

WATER WAVES ABOVE A SILL

YURIKO YAMAMURO

The theory of trapping of water waves in three-dimensional flows, where the bottom topographies model the natural variations of the ocean floors, is difficult (Meyer [6], [7]). Hence, the idealised problem of a submerged circular sill has received the most analysis (Longuet-Higgins [2]; Summerfield [9]; Pite [8]). All these works are based on linear shallow-water equations. In order to verify the large near-resonances predicted by these theories, W.G. Pritchard (Private Communication) performed laboratory experiments and produced results which were inconsistent with the theoretical results. This thesis develops a full linear theory for both inviscid and viscous cases (Havelock [1]; Mahony and Pritchard [3]), and thus serves to define the limitations in the results obtained by using shallow-water theory. The general trend of these results is that wave-trapping is a phenomenon which exhibits less peaky resonance than has been generally assumed. However, Pritchard's experiments do show marked peaks at frequencies not associated with resonances predicted by shallow-water theory.

Since subharmonic resonance can lead to large responses, a weakly nonlinear theory (McEwan [5]; Mahony and Smith [4]), in which the calculations must be organised in a non-standard way, is described. The velocity potential is conveniently written as that of the linear solution plus the components excited by the nonlinear interactions. The calculation of the modal decomposition of the latter is complex and its difficulties are resolved in Chapter 4. A possible subharmonic resonance is presented, together with a particular laboratory condition, under which it can occur.

Received 20 November 1980. Thesis submitted to the University of Western Australia, July 1980. Degree approved, October 1980. Supervisor: Professor J.J. Mahoney.

The response curves exhibit the unusual property that the forced linear mode is amplified as it feeds energy into the half-frequency mode. It is concluded that such interactions are unlikely to have occurred in Pritchard's experiments, and that the inconsistencies in his results may be due to resonances associated with modes of the wave tank.

References

- [1] T.H. Havelock, "Forced surface-waves on water", *Philos. Mag.* (7) 51 (1929), 569-576.
- [2] M.S. Longuet-Higgins, "On the trapping of wave energy round islands", *J. Fluid Mech.* 29 (1967), 781-821.
- [3] J.J. Mahony and W.G. Pritchard, "Wave reflection from beaches" (Fluid Mechanics Research Institute Report 95. University of Essex, 1978).
- [4] J.J. Mahony and Ronald Smith, "On a model representation for certain spatial-resonance phenomena", *J. Fluid Mech.* 53 (1972), 193-207.
- [5] A.D. McEwan, "Degeneration of resonantly-excited standing internal gravity waves", *J. Fluid Mech.* 50 (1971), 431-448.
- [6] R.E. Meyer, "Resonance of unbounded water bodies", *Mathematical problems in the geophysical sciences*. Vol. 1: *Geophysical fluid dynamics*, 189-227 (Lectures in Applied Mathematics, 13. American Mathematical Society, Providence, Rhode Island, 1971).
- [7] R.E. Meyer, "Theory of water-wave refraction", *Adv. Appl. Mech.* 19 (1979), 53-141.
- [8] H.D. Pite, "The excitation of damped waves diffracted over a submerged circular sill", *J. Fluid Mech.* 82 (1977), 621-641.
- [9] W.C. Summerfield, "On the trapping of wave energy by bottom topography" (Horace Lamp Centre Ocean. Res. Paper 30. Flinders University, South Australia, 1969).

Mathematics Research Center,
University of Wisconsin-Madison,
610 Walnut Street,
Wisconsin 53706, USA.